



Gruppen, E. G., Kunutsor, S. K., Kieneker, L. M., van der Vegt, B., Connelly, M. A., de Bock, G. H., Gansevoort, R. T., Bakker, S. J. L., & Dullaart, R. P. F. (2019). GlycA, a novel pro-inflammatory glycoprotein biomarker is associated with mortality: results from the PREVEND study and meta-analysis. *Journal of Internal Medicine*, 286(5), 596-609. <https://doi.org/10.1111/joim.12953>

Publisher's PDF, also known as Version of record

License (if available):
CC BY-NC

Link to published version (if available):
[10.1111/joim.12953](https://doi.org/10.1111/joim.12953)


[Link to publication record in Explore Bristol Research](#)
PDF-document

University of Bristol - Explore Bristol Research

General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available:
<http://www.bristol.ac.uk/red/research-policy/pure/user-guides/ebr-terms/>

Supplemental figure 1



Supplemental Table 1. Sex-stratified analyses between hsCRP and cancer mortality in 5,526 participants (380 events) of the PREVEND study.

	Quartile 1	Quartile 2	P- value	Quartile 3	P-value	Quartile 4	P-value	P for trend*
Men								
Participants (n)	657	665		652		659		
Range, mg/L	<0.62	≥0.62		≥1.31		≥2.84		
No. of deaths (%)	18 (2.7)	49 (7.4)		74 (11.3)		107 (16.2)		
Crude	Ref.	2.74 [1.60-4.70]	<0.001	4.45 [2.66-7.45]	<0.001	6.68 [4.06-11.01]	<0.001	<0.001
Model 1	Ref.	1.85 [1.07-3.19]	0.027	2.50 [1.48-4.25]	0.001	3.11 [1.86-5.22]	<0.001	<0.001
Model 2	Ref.	1.89 [1.10-3.27]	0.022	2.55 [1.50-4.33]	0.001	3.11 [1.85-5.24]	<0.001	<0.001
Model 3	Ref.	1.90 [1.10-3.29]	0.021	2.57 [1.51-4.37]	<0.001	3.06 [1.81-5.16]	<0.001	<0.001
Model 4	Ref.	1.90 [1.10-3.29]	0.022	2.54 [1.49-4.33]	0.001	3.01 [1.78-5.10]	<0.001	<0.001
Model 5	Ref.	1.93 [1.11-3.34]	0.019	2.55 [1.50-4.35]	0.001	2.98 [1.76-5.06]	<0.001	<0.001
Model 6	Ref.	1.86 [1.07-3.22]	0.028	2.37 [1.38-4.08]	0.002	2.57 [1.46-4.53]	0.001	0.019
Women								
Participants (n)	727	719		722		725		
Range, mg/L	<0.63	≥0.63		≥1.41		≥3.25		
No. of deaths (%)	21 (2.9)	32 (4.5)		41 (5.7)		38 (5.2)		
Crude	Ref.	1.52 [0.88-2.63]	0.14	1.95 [1.15-3.30]	0.013	1.84 [1.08-3.13]	0.025	0.08
Model 1	Ref.	1.02 [0.58-1.79]	0.95	1.13 [0.65-1.98]	0.66	1.09 [0.61-1.95]	0.78	0.82
Model 2	Ref.	1.04 [0.59-1.83]	0.89	1.09 [0.62-1.90]	0.76	1.07 [0.59-1.92]	0.83	0.90
Model 3	Ref.	1.09 [0.62-1.93]	0.77	1.15 [0.65-2.05]	0.63	1.12 [0.61-2.05]	0.71	0.85
Model 4	Ref.	1.04 [0.59-1.85]	0.89	1.12 [0.63-2.00]	0.70	1.07 [0.59-1.96]	0.82	0.91
Model 5	Ref.	1.04 [0.59-1.86]	0.86	1.12 [0.62-2.00]	0.71	1.08 [0.58-1.98]	0.82	0.90
Model 6	Ref.	1.01 [0.57-1.81]	0.96	1.06 [0.58-1.92]	0.86	0.95 [0.481-1.90]	0.88	0.77

Hazard ratios were derived from Cox proportional hazards regression models.

Model 1: crude model + age, sex, BMI, alcohol intake (<10g/d or >10 g/d) and smoking status (never, former current).

Model 2: model 1 + diabetes, systolic blood pressure, lipid lowering drugs and anti-hypertensive medications.

Model 3: model 2 + total cholesterol, HDL cholesterol and triglycerides.

Model 4: Model 3 + history of CVD and history of cancer.

Model 5: Model 4 + eGFR_{creatinine} cystatin C and UAE.

Model 6: Model 5 + GlycA

Triglycerides, UAE and hsCRP were log transformed when used as a continuous variable in the analyses.

*Tests of trend across increasing quartiles were conducted by assigning the median for each quartile as its value and treating this as a continuous variable.

Abbreviations: *BMI*, body mass index; *HDL-cholesterol*, high density lipoprotein cholesterol; *CVD*, cardiovascular disease; *hsCRP*, high-sensitivity C-reactive protein; *UAE*, urinary albumin excretion; *PREVEND*, Prevention of REnal and Vascular ENd-stage Disease.

Supplemental Table 2. Sex-stratified analyses between GlycA and cancer mortality in 5,526 participants (380 events) of the PREVEND study.

	Quartile 1	Quartile 2	P- value	Quartile 3	P-value	Quartile 4	P-value	P for trend*
Men								
Participants (n)	655	644		667		667		
Range, $\mu\text{mol/L}$	<305	≥ 305		≥ 338		≥ 381		
No. of deaths (%)	30 (4.6)	47 (7.3)		70 (10.5)		101 (15.1)		
Crude	Ref.	1.64 [1.04-2.60]	0.034	2.41 [1.57-3.70]	<0.001	3.72 [2.48-5.59]	<0.001	<0.001
Model 1	Ref.	1.04 [0.65-1.66]	0.19	1.34 [0.86-2.06]	0.19	1.82 [1.19-2.79]	0.006	<0.001
Model 2	Ref.	1.02 [0.64-1.63]	0.94	1.32 [0.85-2.05]	0.21	1.78 [1.16-2.74]	0.009	<0.001
Model 3	Ref.	1.06 [0.66-1.70]	0.81	1.37 [0.87-2.13]	0.17	1.87 [1.20-2.90]	0.006	<0.001
Model 4	Ref.	1.07 [0.67-1.72]	0.78	1.36 [0.87-2.12]	0.18	1.84 [1.18-2.87]	0.007	0.001
Model 5	Ref.	1.08 [0.67-1.73]	0.76	1.35 [0.86-2.11]	0.19	1.82 [1.16-2.84]	0.009	0.001
Model 6	Ref.	0.99 [0.62-1.60]	0.98	1.14 [0.71-1.81]	0.59	1.35 [0.82-2.22]	0.24	0.12
Women								
Participants (n)	712	715		743		723		
Range, $\mu\text{mol/L}$	<313	≥ 313		≥ 352		≥ 394		
No. of deaths (%)	18 (2.5)	27 (3.8)		45 (6.1)		42 (5.8)		
Crude	Ref.	1.49 [0.82-2.70]	0.19	2.43 [1.41-4.20]	0.001	2.35 [1.35-4.07]	0.002	0.001
Model 1	Ref.	1.06 [0.58-1.94]	0.86	1.49 [0.84-2.64]	0.17	1.38 [0.76-2.49]	0.29	0.20
Model 2	Ref.	1.05 [0.57-1.92]	0.88	1.39 [0.78-2.47]	0.26	1.29 [0.71-2.34]	0.41	0.33
Model 3	Ref.	1.09 [0.59-2.00]	0.79	1.40 [0.78-2.52]	0.27	1.36 [0.74-2.52]	0.32	0.27
Model 4	Ref.	1.09 [0.59-2.00]	0.79	1.47 [0.81-2.64]	0.20	1.40 [0.76-2.58]	0.28	0.22
Model 5	Ref.	1.10 [0.60-2.02]	0.77	1.47 [0.82-2.65]	0.20	1.41 [0.76-2.60]	0.27	0.22
Model 6	Ref.	1.11 [0.60-2.06]	0.73	1.53 [0.83-2.81]	0.18	1.51 [0.76-3.01]	0.24	0.19

Hazard ratios were derived from Cox proportional hazards regression models.

Model 1: crude model + age, sex, BMI, alcohol intake (<10g/d or >10 g/d) and smoking status (never, former current).

Model 2: model 1 + diabetes, systolic blood pressure, lipid lowering drugs and anti-hypertensive medications.

Model 3: model 2 + total cholesterol, HDL cholesterol and triglycerides.

Model 4: Model 3 + history of CVD and history of cancer.

Model 5: Model 4 + eGFR_{creatinine cystatin C} and UAE.

Model 6: Model 5 + hsCRP

Triglycerides, UAE and hsCRP were log transformed when used as a continuous variable in the analyses.

*Tests of trend across increasing quartiles were conducted by assigning the median for each quartile as its value and treating this as a continuous variable.

Abbreviations: *BMI*, body mass index; *HDL-cholesterol*, high density lipoprotein cholesterol; *CVD*, cardiovascular disease; *hsCRP*, high-sensitivity C-reactive protein; *UAE*, urinary albumin excretion; *PREVEND*, Prevention of Renal and Vascular End-stage Disease.

Supplemental Table 3. Association between GlycA and hsCRP levels and lung cancer mortality in 5,526 participants (57 events) of the PREVEND study.

	Per 1 SD increment	P- value
GlycA		
Crude	1.61 [1.33-1.95]	<0.001
Model 1	1.34 [1.07-1.67]	0.011
Model 2	1.33 [1.06-1.67]	0.015
Model 3	1.34 [1.07-1.67]	0.011
Model 4	1.38 [1.02-1.88]	0.040
hsCRP		
	Per 1 SD increment	
Crude	1.48 [1.19-1.84]	<0.001
Model 1	1.18 [0.92-1.52]	0.19
Model 2	1.17 [0.91-1.50]	0.23
Model 3	1.18 [0.92-1.51]	0.19
Model 4	0.96 [0.70-1.33]	0.81

Hazard Ratios are given per 1 SD increment in GlycA and hsCRP

Model 1: crude model + age, sex, BMI, alcohol intake (<10g/d or >10 g/d) and smoking status (never, former current).

Model 2: model 1 + history of CVD and history of cancer.

Model 3: model 2 + eGFR_{creatinine cystatin C} and UAE.

Model 4: Model 3 + hsCRP (for GlycA analyses) + GlycA (for hsCRP analyses).

UAE and hsCRP were log transformed when used as a continuous variable in the analyses.

1 SD is 62 µmol/L for GlycA and 5.22 for hsCRP. Abbreviations: *BMI*, body mass index; *CVD*, cardiovascular disease; *hsCRP*, high-sensitivity C-reactive protein; *UAE*, urinary albumin excretion; *PREVEND*, Prevention of RENal and Vascular ENd-stage Disease.

Supplemental Table 4. Association between GlycA and hsCRP levels and **gastrointestinal cancer mortality** in 5,526 participants (56 events) of the PREVENT study.

	Per 1 SD increment	P- value
GlycA		
Crude	1.31 [1.04-1.65]	0.023
Model 1	1.39 [1.09-1.75]	0.007
Model 2	1.40 [1.10-1.79]	0.007
Model 3	1.42 [1.11-1.83]	0.006
Model 4	1.41 [1.01-1.98]	0.043
hsCRP		
Crude	1.22 [0.97-1.55]	0.094
Model 1	1.25 [0.97-1.61]	0.08
Model 2	1.24 [0.97-1.60]	0.09
Model 3	1.25 [0.97-1.61]	0.08
Model 4	1.01 [0.73-1.41]	0.95

Hazard Ratios are given per 1 SD increment in GlycA and hsCRP

The following organs belong to the category of gastrointestinal: bile duct, esophagus, colon + rectum, stomach, pancreas, small intestine and liver.

Model 1: crude model + age, sex, BMI, alcohol intake (<10g/d or >10 g/d) and smoking status (never, former current).

Model 2: model 1 + history of CVD and history of cancer.

Model 3: model 2 + eGFR_{creatinine cystatin C} and UAE.

Model 4: Model 3 + hsCRP (for GlycA analyses) + GlycA (for hsCRP analyses).

UAE and hsCRP were log transformed when used as a continuous variable in the analyses.

1 SD is 62 µmol/L for GlycA and 5.22 mg/L for hsCRP. Abbreviations: *BMI*, body mass index; *CVD*, cardiovascular disease; *hsCRP*, high-sensitivity C-reactive protein; *UAE*, urinary albumin excretion; *PREVENT*, Prevention of RENal and Vascular ENd-stage Disease.